

MFC Injection

For animal use only

Reg.no: G329 (Act 36/1947)

Indications for use:

For the treatment of milk fever in cattle and sheep, especially where this is complicated by low blood levels of magnesium and/or phosphorus.

Concurrent acetonemia is also controlled by the inclusion of dextrose. This also meets increased sugar requirements in cases of milk fever.

Composition:

Calcium borogluconate 20.9 % m/v
Magnesium hypophosphite 4.2 % m/v
Dextrose 16.7 % m/v

Warnings:

- Keep out of reach of children, uninformed persons and animals.
- Pain and transient local tissue swelling may sometimes follow subcutaneous injections especially in thin skinned subjects. It is, therefore, recommended that the intravenous route be used whenever possible.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide variety of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

Directions for use: Use only as directed

The remedy must be warmed to body temperature prior to use by placing the bottle in warm water. Intravenous injection should be carried out slowly.

Cattle: 200 - 400 ml dose

Sheep: 25 - 75 ml dose

Storage:

Store in a cool, dry place below 25°C.
Protect from direct light.

Registration holder:

Lionel's Veterinary Supplies (Pty) Ltd
2001/011520/07
PO Box 901, Sanlamhof, 7532
Tel. 021 932 2019
www.lionelsvet.co.za

