

CBG Injection

Calcium Borogluconate Injection - For animal use only

Reg.no: G1095 (Act 36/1947)

Indications for use:

For the treatment of milk fever in cows and lambing sickness in ewes.

Composition:

Calcium gluconate 33.33 % m/v

Boric acid 6.67 % m/v

Calcium borogluconate is available as a sterile 40 % m/v aqueous solution, containing 0.1 % chlorocresol as a preservative.

Warnings:

- Keep out of reach of children, uninformed persons and animals.
- Failure in cases of milk fever may be due to a concomitant hypomagnesaemia and/or hypophosphataemia or even to acetoanaemia - cases where your veterinarian should be consulted.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide variety of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

Directions for use: Use only as directed

By subcutaneous or intravenous injection.

The remedy must be warmed (to approximately 37°C) prior to use and if injected intravenously, it must be given slowly.

Cows: 210 - 400 ml dose (84 - 160 g approved CBG content)

Sheep and goats: 35 ml dose (14 g approved CBG content)

Storage:

Store in a cool, dry place below 25°C.
Protect from direct light.

Registration holder:

Lionel's Veterinary Supplies (Pty) Ltd
2001/011520/07
PO Box 901, Sanlamhof, 7532
Tel. 021 932 2019
www.lionelsvet.co.za

