



bontera[®]

Richer earth. Better harvest.

IMPACT

Organic Microbial Soil Enhancer

Water-Soluble Powder Concentrate

Registration number M34, Act 36 of 1947 FERTILIZER GROUP 3

Product suitable for use in Organic Farming in accordance with EU Regulations 2018/848 and 2021/1165.

Registered plant nutrient content:



Humic Acid derived from Leonardite	38.14%
Fulvic Acid	49.10%
Humins	8.25%
Microbes (Bacteria and Fungi)	3.10%
Other Ingredients	1.40%

pH 7.2 - 7.6 *Bacillus spp.* 2.0 x 10⁷ cfu/g *Trichoderma spp.* 3.0 x 10⁶ cfu/g



**Controlled By
ECOCERT F-32600**



**Bontera is suited for a wide variety of applications, including:
Agriculture, Horticulture, Floriculture, Orchards and Pastures**

Precautionary Statements:

Read carefully and follow all instructions.
Keep out of reach of children.
If medical advice is needed, have a product container or label at hand.

REGISTRATION HOLDER: Bontera SA (Pty) Ltd
Company Registration Number: 2013/077999/07
Physical Address: Suite 2-102, Monument Office Park, 71 Steenbok Ave, Monument Park, Pretoria, 0181 Telephone number: +27 12 347 0584

1 kg

Net Contents: 1kg / 2.2 lbs.
10 units per case

DIRECTIONS FOR USE:

Use only as directed. One 1kg bag is equal to 10 Litres of liquid concentrate. Dosage below per litre is based on the amount of liquid concentrate required. Ensure that mixing tanks and equipment are free of chemical residues and use fresh, clean water for the dilution. It is preferable to apply the product in the late afternoon or in cooler weather, avoiding UV-rays and extreme temperatures. Use mixed product within 18 hours.

SHAKE WELL BEFORE USE!

SOIL APPLICATION:

- 1) 1-1.5 L/ha (11-21 fl. oz/acre) at planting;
- 2) 1-3 applications of 0.8-1.0 L/ha (11-14 fl. oz/ acre) at appropriate growth stages depending on plant/ crop growth cycle. Use drip irrigation or other irrigation systems to ensure the product is well dispersed over the entire area.

FOLIAR APPLICATION:

Spray 1:800 dilution of the liquid concentrate until foliage is completely wet at (1) new leaf emergence; (2) flower bud initiation; (3) re-growth; and (4) any growth stage thereafter that is considered important depending on plant/crop growth cycle. For specific application guidelines, please consult your agent.

SEED SOAKING (OPTIONAL)

Soak seeds for 30min in 1:300 dilution of the liquid concentrate prior to planting. The dose of 1st soil application can be reduced by half.

TRANSPLANTS:

Apply 1:10,000 dilution of the liquid concentrate directly into the transplanting hole around the roots until the soil is wet.

For specific application guidelines, please consult your agent


CO-APPLICATION: **Bontera® IMPACT** can be co-applied with chemical fertilizers. DO NOT APPLY **Bontera® IMPACT** with fungicides.

STORAGE: Store in an airtight container under dry/cool conditions (5-30°C;40- 85°F). Avoid exposure to direct sunlight.

DISCLAIMER: Recommendations for the use of this product are based upon tests deemed to be reliable. The use of this product is beyond the control of the manufacturer. No guarantee, expressed or implied, is made as to the effects of such or the results to be obtained. The buyer assumes all responsibility, including injury or damage resulting from its misuse, including its application in combination with other products.

Batch No / Manufacturing Date / Expiry Date

BONWP2



www.bontera.co.za