



bontera[®]

Richer earth. Better harvest.

IMPACT XTRA

Microbial Soil Enhancer

Water-Soluble Powder Concentrate

Registration number M155, Act 36 of 1947 FERTILIZER GROUP 3

Registered plant nutrient content:

A blend of assorted microbial cultures in humic and fulvic acids

Humic Acid 189.7 g/kg, Fulvic Acid 674.6 g/kg
Bacillus amyloliquefaciens 1.5×10^8 cfu/gram
Bacillus pumilus 3.0×10^8 cfu/gram
Bacillus subtilis 3.5×10^7 cfu/gram
Bacillus licheniformis 5.5×10^7 cfu/gram
Bacillus megaterium 3.0×10^7 cfu/gram
Trichoderma harzianum 1.0×10^9 cfu/gram
Trichoderma koningii 2.0×10^6 cfu/gram



Precautionary Statements:

Read carefully and follow all instructions.
Keep out of reach of children.
If medical advice is needed, have a product container or label at hand.

REGISTRATION HOLDER: Bontera SA (Pty) Ltd
(Reg no: 2013/077999/07) Suite 2-102, Monument
Office Park, 71 Steenbok Avenue, Monument
Park, 0181, +27 12 347 0584 info@bontera.co.za
Made in USA

1 kg

Net Contents: 1kg / 2.2 lbs.
10 units per case

Bontera® Impact Xtra is suited to a wide variety of applications, including Agricultural and Horticultural crops.

Bontera® Impact Xtra contains a unique combination of beneficial microbes, humic and fulvic acids intended for use as a microbial soil enhancer. It improves the growth and/or yield of plants.

DIRECTIONS FOR USE: Use only as directed.

Crop	Application Instructions
Bulb & Leafy Vegetables Broccoli, Brussel sprout, Cabbage, Garlic, Leeks, Lettuce, Spinach, Onion, etc. Solanaceous Fruits Tomato, Green pepper, Paprika, Chilli, etc. Cucurbits Butternut, Gem squash, Cucumber, Baby marrow, Patty pan, etc.	i) Apply 100 - 150 g/ha as a soil drench at planting ii) Apply 1-3 applications of 80 - 100 g/ha at appropriate growth stages depending on plant/crop growth cycle. Use drip irrigation or other irrigation systems to ensure the product is well dispersed over the entire area.
Row Crops: Grains Maize, Sweetcorn, Sorghum, Millet, etc.	Apply 100 - 200 g/ha as a soil drench at planting, with a follow-up at four weeks after planting.
Potato All varieties	Apply 200 g/ha at planting as a soil drench with a follow-up at six and ten weeks after planting.
Tree Crops: Citrus Orange, Lemon, Mandarin, etc.	Apply 100 - 200 g/ha as a soil drench two to three times during the growing season.

CO-APPLICATION:

Bontera® Impact Xtra can be co-applied with chemical fertilisers. **DO NOT APPLY Bontera® Impact Xtra** with fungicides or nematicides.

STORAGE: Store in an airtight container under dry/cool conditions (5-30°C). Avoid exposure to direct sunlight.

DISCLAIMER: Recommendations for the use of this product are based upon tests deemed to be reliable. The use of this product is beyond the control of the manufacturer. No guarantee, expressed or implied, is made as to the effects of such or the results to be obtained. The buyer assumes all responsibility, including injury or damage resulting from its misuse, including its application in combination with other products. For all other crops not listed on this label the user should determine the suitability of the product for the intended use and user assumes the risk and liability in connection therewith. The user does not have the right to de-formulate this product, or to isolate and/or culture any of the active ingredient for any purpose. The customer is not permitted to re-sell the product in any form without the prior written authorisation from the Registration Holder.

Batch No / Manufacturing Date / Expiry Date

BONIX2



www.bontera.co.za