

MAXICARE®



For animal use only
 Reg.no: G3145 (Act 36/1947)
 Namibia Reg. no: V02/18.1.8/762 (Act 13/2003)

A roundworm, lungworm, milk tapeworm, liver fluke and nasal bot remedy with residual protection against re-infestation of wireworm and hookworm in sheep and goats

Composition:

Albendazole 1,9 % m/v
 Closantel 3,0 % m/v

Direction for use: Use only as directed

Use as an oral drench. Shake well before use.
 Dose rate for sheep and goats: 2,0 ml / 10 kg body mass.

Body mass (kg)	Dosage (ml)	Body mass (kg)	Dosage (ml)
10	2	40	8
20	4	50	10
30	6		

Warnings:

NOT FOR USE IN MILK GOATS WHOSE MILK IS INTENDED FOR HUMAN CONSUMPTION.
 Allow 30 days between last dosing and slaughter for human consumption.
 Handle with care.
 Management practices should always ensure that all drenching procedures are carefully carried out. It is particularly important not to exceed the recommended dose in ewes during the mating season, or during the first month of pregnancy.
 Keep out of reach of uninformed persons, children and animals.
 Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

Precautions:

- It is recommended that all sheep be vaccinated against Pulpy Kidney Disease before any worm remedy is administered.
- Do not treat animals more frequently than every three weeks.
- Wash hands thoroughly after use.

Storage:

Store in cool place and protect from sunlight.

Registration holder:

CEVA Animal Health (Pty) Ltd (Reg. No. 1973/016009/07), P O Box 7707, Halfway House 1685
 011 312 4088

Efficacy:

Worm species	Adult	Immatures
Wireworm	*	*
Brown stomachworm	*	*
Bankruptworm	*	*
Long-necked bankruptworm	*	*
Hookworm	*	*
Nodular worm	*	*
Large-mouthed bowelworm	*	*
Liver fluke	*	** (6 weeks old)
Giant liver fluke	*	** (8 weeks old)
Milk tapeworm	Class 1	

*Controls 90 - 100 % effective

**Aids in control = 60 - 89 % effective

Class 1 = 100 % effective in more than 80 % of treated animals. Controls all stages of nasal bot. Prevents roundworm eggs present in animal at time of dosing, from hatching.

Residual effect:

Worm species	Interval between treatment and re-infestation
Wire worm	3 weeks
Hook worm	3 weeks

