



PRODUCT DATASHEET

GENERAL

For the treatment of secondary bacterial infections in poultry sensitive to sulphamethoxy pyridazine and trimethoprim.

MODE OF ACTION

The trimethoprim phenyl-propanol-disulphonate sodium is a potentiator of sulphonamide, which after oral ingestion is completely and rapidly absorbed within 1 to 5 hours. Maximum serum levels are reached 2 to 4 hours after oral ingestion. Trimethoprim phenyl-propanol-disulphonate sodium is widely distributed throughout the body tissues and can be found in the kidney, liver and lungs.

COMPOSITION

Sulphamethoxy pyridazine 69%
Trimethoprim 21%.

SPECIFICATION

Colour: White / Yellow Powder.

DOSAGE AND DIRECTIONS FOR USE

Use only as directed: Oral administration only.

HOUSE TYPE	DOSAGE (G) AND ADMINISTRATION IN DRINKING WATER
Convection Housing	22 grams of Coli-Mix Plus® in 100 litres of drinking water for 5 consecutive days. In severe infections, the dosage may be doubled to 44 grams per 100 litres drinking water for the first day of treatment only.
Controlled Environmental Housing	100 grams of Coli-Mix Plus® per 100 litres of drinking water in 4- to 5-week-old broilers for three days running followed by 50 grams per 100 litres drinking water for 2 - 3 days thereafter.

WARNINGS

Withdrawal period: Meat and Eggs - 72 hours

Do not use for layer hens: for use in broilers and pullets only.

Not for use in humans.

Keep out of reach of children, uninformed persons and animals. Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide variety of reasons. If this is suspected, seek veterinary advice, and notify the registration holder.

PRECAUTIONS

Symptoms of toxicity are more likely to occur with the prolonged administration of high doses.

STORAGE

Store in a cool, dry place.

PACKAGING

100 grams, 10 kilograms and 25 kilograms.

